Current Treatment Strategies for Patella Fractures

Dustin J. Schuett, DO; Mark E. Hake, MD; Cyril Mauffrey, MD, FACS, FRCS; E. Mark Hammerberg, MD; Philip F. Stahel, MD; David J. Hak, MD, MBA

Abstract: Patella fractures can be caused by excessive tension through the extensor mechanism or a direct impact. Nondisplaced fractures with an intact extensor mechanism can be treated nonoperatively. Surgical treatment is recommended for fractures that either disrupt the extensor mechanism or have greater than 2 to 3 mm of step-off and greater than 1 to 4 mm of displacement. Tension band fixation is the most commonly employed surgical technique; however, this can be technically demanding, especially in multifragmentary fractures. Symptomatic hardware is the most common complication following operative treatment. Functional impairment remains common after treatment of patella fractures. The purpose of this article is to review current treatment strategies to help optimize the management of patients with such patella fractures. [*Orthopedics.* 2015; 38(6):377-384.]

Patella fractures account for approximately 1% of frac-

tures in adults.^{1,2} The patella is the largest sesamoid in the

The authors are from the Department of Orthopaedic Surgery (DJS), Naval Medical Center San Diego, San Diego, California; and the Department of Orthopaedic Surgery (MEH, CM, EMH, PFS, DJH), Denver Health/University of Colorado, Denver, Colorado.

Drs Schuett, Hake, and Stahel have no relevant financial relationships to disclose. Dr Mauffrey has received a teaching honorarium from AO, DePuy Synthes, and Abbott Medical. Dr Hammerberg has received study funding from Zimmer. Dr Hak has received personal fees from Invibio, Merck, Kuros Biosurgery, and RTI Biologics.

Correspondence should be addressed to: David J. Hak, MD, MBA, Department of Orthopaedic Surgery, Denver Health/University of Colorado, 777 Bannock St, MC 0188, Denver, CO 80204 (david.hak@dhha.org). doi: 10.3928/01477447-20150603-05 body and functions to increase the moment arm of the extensor mechanism of the quadriceps by 30%.³ The cartilage on the articular surface of the patella is among the thickest in the human body, measuring up to 5.5 mm.^{2,4} Treatment goals for patella fractures include restoring the function of the extensor mechanism, minimizing patellar bone loss, maximizing articular congruity, and allowing early mobilization.^{2,5,6}

Patella fractures can occur as a result of either an indirect or a direct mechanism. The indirect mechanism typically results in a transverse fracture, produced when the force exerted by the extensor mechanism exceeds the tensile strength of the patella.⁷ A direct blow to the anterior patella, typically with the knee flexed, causes failure of the patella in compression, often resulting in a comminuted fracture.⁷

Comminuted fractures comprise 55% of surgically treated patella fractures.^{1,6} Operative treatment of comminuted patella fractures presents a significant challenge to surgeons.⁸ Although some comminuted fractures with an intact extensor mechanism can be treated nonoperatively, surgical treatment is indicated for extensor mechanism disruption, greater than 2 to 3 mm of articular step-off, and greater than 1 to 4 mm of displacement.^{1,2} Failure to restore the articular surface contour results in posttraumatic arthritis.⁹

Unfortunately, functional impairment remains common after treatment of patella fractures. Functional outcome measures were prospectively collected at 3, 6, and 12 months following surgical fixation of unilateral patella fractures in 30 patients. Twenty-four patients (80%) experienced anterior knee pain during activities of daily living. At 12 months, objective testing demonstrated that, compared with the uninjured knee, the knee extensor mechanism on the injured side had an average 41% deficit in strength, 47% deficit in power, and 34% deficit in endurance.10 In another study, involving 40 patients who underwent surgical treatment of an isolated, unilateral patella fracture with an average follow-up of 6.5 years (range, 1.25-17 years), investigators found that significant symptomatic complaints and functional deficits persisted based on validated outcome measures as well as objective physical evaluations.11 Removal of symptomatic hardware was performed in 52% of the patients treated with osteosynthesis, whereas 38% of the patients with retained hardware reported implant-related pain at least some of the time. Eight patients (20%) had an extensor lag greater than 5°, 15 patients (38%) had a restricted range of flexion greater than 5°, and 6 patients (15%) had a restricted range of extension greater than 5°. Patients had an average isometric extension deficit of 26% between the uninvolved and involved sides for peak torque. Mean normalized SF-36 physical composite score and mean normalized Knee Injury and Osteoarthritis Outcome Scores subscale scores were statistically different (P<.05) from reference population norms.

IMAGING

Typically, fracture classification and treatment decisions are based on quality anteroposterior and lateral radiographs of the knee, with advanced imaging rarely being indicated.² However, a recent study by Lazaro et al¹² found that classification according to the AO/OTA system was modified in 66% of cases and treatment plans were changed for 49% of patients after obtaining computed tomography scans. Most commonly, they identified a severely comminuted distal pole fracture, which was missed on nearly half of plain films.

TREATMENT Nonoperative Treatment

Nonoperative management consisting of immobilization for 4 weeks results in good to excellent results in 99% of patients with an intact extensor mechanism, less than 4 mm of displacement, and less than 3 mm of articular stepoff.1,2 Prolonged immobilization can result in knee joint stiffness, quadriceps atrophy, and joint adhesions.9 More recently, Melvin and Mehta² recommended weight bearing as tolerated in a knee immobilizer locked in extension with straight-leg raises when tolerated followed by active and active assist range of motion at 1 to 2 weeks and resistance activities at 6 weeks.

Nonoperative management should be considered when significant patient comorbidities make operative intervention perilous.² Pritchett¹³ reported on 18 patella fractures in medically complex patients with more than 1 cm of displacement managed with a hook and loop fastener splint, full weight bearing, and straight-leg raises. Six patients had died by the 24-month follow-up, and 3 of the surviving 12 patients had poor outcomes.

Operative Treatment

Numerous treatment methods, including partial excision, tension band, modified tension band, osteosynthesis with plates and screws, suture repair, cerclage wiring, percutaneous open reduction and internal fixation, total patellectomy, arthroscopic-assisted open reduction and internal fixation, and external fixation, have been used for patella fractures.^{2,5,6,14-16} A recent review found little high-quality evidence comparing operative treatment modalities for patella fractures and a similar paucity of evidence comparing operative with nonoperative treatment.17

Tension Band Fixation

For simple transverse fractures, the most widely used technique is open reduction and internal fixation with an anterior tension band (Figure 1).^{5,16} This technique is designed to convert the tension force seen by the anterior patella into a compressive force across the articular surface, promoting fracture healing.^{5,18} Although tension band fixation per the classic AO technique is reserved for simple fractures, certain comminuted fractures can also be treated with a tension band construct if the posterior cortex is intact to allow for compression.¹⁸

Surgical fixation is typically performed through an anterior longitudinal incision augmented as needed by fullthickness medial and lateral flaps.⁵ Some authors have advocated a lateral parapatellar approach to allow improved exposure of the fracture for reduction and fixation; however, a transverse approach should not be used unless an open wound dictates otherwise.^{19,20} Tension band fixation allows early motion, which has been found to improve outcomes and decrease posttraumatic arthritis.^{18,20,21}

The classic AO technique described in the 1950s consists of 2 parallel vertical Kirschner wires with a tension band passing anteriorly over the patella and posterior to the Kirschner wires.18,20 Use of a large-bore plastic catheter can be helpful in passing the tension band wire posterior to the vertical Kirschner wires. This technique has been found to be associated with prominent implants requiring hardware removal, implant migration, muscular atrophy, and loss of reduction.7,21

Berg²¹ reported that a modified tension band technique using parallel vertical cannulated screws resulted in union in all 10 patients, including 3 revision cases, with 70% good to excellent outcomes and no loss of reduction, implant migration, or implant failure. Tian et al²² performed a retrospective review comparing a modified tension band technique using Kirschner wires with a modified tension band technique using cannulated screws. They found improved fracture reduction, a reduced healing score, and better Iowa knee scores with the cannulated screw modification. Additionally, the implant migration rate and the second operation rate were 15.4% and 5.7%,



Figure 1: Anteroposterior (A) and lateral (B) radiographs of a transverse patella fracture amenable to tension band wiring. Anteroposterior (C) and lateral (D) views showing healing of the fracture 5 months postoperatively.

respectively, for the Kirschner wire, with no complications in the cannulated screw cohort. In a cadaveric study, Carpenter et al^{23} found that a modified tension band with cannulated screws had a higher load to failure than cannulated screws alone or a modified tension band with Kirschner wires.

Gosal et al^{24} compared stainless steel wire with braided polyester using the Kirschner wire tension band technique. The reoperation rate for the steel wire group was 38%, compared with 6% for the braided polyester group. The current recommendation for transverse patella fractures is figure-of-8 tension band wiring with vertical cannulated screws.²

Cerclage Wire Fixation

Yang et al²⁵ described a titanium cable cerclage wiring technique that they used to treat displaced comminuted patella fractures in 21 patients. They had 1 case of cerclage wire breakage and 100% good to excellent results with 100% union rates. Matsuo et al²⁶ performed cerclage wire fixation including the surrounding soft tissue in their repair of



Figure 2: Anteroposterior (A) and lateral (B) radiographs of a comminuted patella fracture. Open reduction was performed, followed by placement of 2.0and 2.7-mm screws. A 2.7-mm plate was added for increased stability. Anteroposterior (C) and lateral (D) radiographs showing maintenance of reduction and healing of the fracture.

5 patients with comminuted fractures and reported an 80% union rate with 1 inferior pole nonunion and no extensor lag. They stated that incorporating the soft tissues into their repair permitted the use of this technique for comminuted fractures not amenable to tension band fixation.

Plate Fixation

Small plates can be applied to the anterior surface of the patella in the setting of comminution to provide additional stability (**Figure 2**). Taylor et al²⁷ recently reported techniques and outcomes of plate fixation for patella fractures. They presented 8 patients with patella fractures or nonunions treated with a combination of plate and interfragmentary screw fixation. All of their patients went on to union at a mean of 3.2 months with an average total arc of knee motion of 129°. There were no cases of hardware removal for symptomatic implants.

A 2.7-mm, fixed-angle plating construct was evaluated by Thelen et al²⁸ for transverse pa-



tella fractures in cadavers and compared with Kirschner wire tension band fixation and cannulated screw tension band fixation. After 100 cycles of full extension to 90° of flexion, the fixed-angle plating group averaged less than 1 mm of displacement, compared with 7.1 and 3.7 mm in the Kirschner wire and cannulated screw tension band groups, respectively. Banks et al²⁹ compared the tension band construct with cannulated screws with a tension band construct with a locking plate in a cadaveric transverse patella fracture model. The locked plate tension band models had similar load to failure, significantly higher ultimate fixation strength, and slightly lower stiffness at final loading

compared with the cannulated screw tension band construct. In a foam patella model, Wurm et al³⁰ loaded simulated patella fractures fixed with locked plating or tension band fixation to failure in a simulated walking test. They found that the tension band construct had a 33% lower load to failure and 5 times larger fracture gap displacement than the locking plate construct.

The use of a titanium mesh, used in craniomaxillofacial surgery as a buttress for fracture fixation, has also been proposed for patella fracture fixation (**Figure 3**). Benefits of this implant include that it is easily contoured, has multiple holes for screw placement, and is very low profile. In a bio-



Figure 3: Anteroposterior (A) and lateral (B) radiographs of a comminuted patella fracture. Postoperative anteroposterior (C) and lateral (D) radiographs and intraoperative photograph (E) showing the use of titanium mesh for fixation.

mechanical study, Dickens et al³¹ found that, compared with standard tension band fixation, the titanium mesh fixation construct maintained a smaller fracture gap prior to failure.³¹

Isolated Interfragmentary Screw Fixation

Little has been published in the literature on isolated screw fixation for patella fractures. Wang et al³² published a retrospective review of transverse patella fractures involving 37 patients treated with modified tension bands compared with 35 patients treated with parallel interfragmentary screws in a lag by design fashion. They found parallel titanium screw fixation to have a shorter operative time, a lower loss of fixation, and lower rates of symptomatic hardware and second surgery.

Tandogan et al⁹ reported on 5 patients with displaced patella fractures without extensor mechanism disruption treated with arthroscopicassisted reduction and percutaneous screw fixation. Their technique returned all but 1 patient to full range of motion with no implant failure or infection. Cadaveric studies have demonstrated that cannulated screws have a lower load to failure than cannulated screws with a modified tension band technique.²³

Inferior Pole Fracture Treatment

Avulsion fractures of the inferior patella account for 9% to 22% of surgically treated patella fractures. These fractures are often comminuted, making treatment difficult.33 Kastelec and Veselko³³ compared the results for 14 patients who had internal fixation of an inferior patella fracture with a basket plate with 14 patients who were treated for the same injury with patellar pole excision and direct patellar tendon repair. At an average of 4.6 years postoperatively, the basket plate group had significantly less pain, a higher activity level, and better knee range of motion than the patients who underwent excision and patellar tendon repair. Patella baja was seen in all but 3 patients in the excision and repair group and was associated with poor functional outcomes. Separate vertical wiring specifically oriented for inferior pole fractures resulted in 100% union in one study of 25 patients with inferior pole patella fractures.34 Partial patellectomy is typically best reserved for comminuted inferior pole fractures not amenable to

fixation.² Egol et al³⁵ compared a cohort of 13 patients with displaced distal pole patella fractures treated with partial patellectomy with a control cohort with central patella fractures treated with tension band techniques. At 1 year, there were no differences between the 2 groups except for more symptomatic hardware requiring intervention in the tension band group. They concluded that suture fixation of distal pole patella fractures is an acceptable technique that yields outcomes similar to tension band treatment of patella fractures.

Partial Patellectomy

A partial patellectomy is performed by first excising comminuted bone fragments and then passing nonabsorbable braided suture from the patellar tendon through drill holes in the patella, similar to a traditional patellar tendon repair (Figure 4). Bone fragments can often be incorporated into the repair. Partial patellectomy has been described as a means to preserve the moment arm of the patella resulting in less loss of strength, ligament instability, and quadriceps atrophy when compared with total patellectomy.36 Bonnaig et al37 compared 26 patients who underwent patella open reduction and internal fixation with 26 patients who underwent partial patellectomy and found no difference in outcomes between the 2 groups. In their study, partial patellectomy was only performed when the treating surgeon felt that an anatomic reduction was not possible.

Minimally Invasive and Percutaneous Techniques

Percutaneous treatment of patella fractures has been proposed as a means to preserve vascular supply and to decrease insult to the soft tissue envelope. In a randomized, controlled trial of 53 patients, Luna-Pizarro et al³⁸ compared the percutaneous patellar osteosynthesis system technique with open surgery for operative patella fractures. They found that the percutaneous patellar osteosynthesis system resulted in shorter surgical time, less pain, better range of motion, fewer complications, and similar functional scores at 2 years postoperatively.

A minimally invasive technique for tension band fixation of transverse patella fractures using the cable pin system (Cable-Ready, Zimmer, Warsaw, Indiana) was evaluated by Mao et al.³⁹ A total of 31 patients were followed for an average of 21 months. Fracture union occurred at a mean of 7.2 weeks with an average of 91° of active flexion at that time. Full range of motion was achieved in 93.5% of the patients at final follow-up and excellent results in 30 of 31 patients.

External Fixation

Wardak et al¹⁴ used a compressive external fixation system for the treatment of 84 displaced primarily transverse patella fractures, of which 31% were open fractures, in Afghanistan. The device was left in place for a total of 6 weeks on average and then removed in a clinic, at which time all fractures had attained



Figure 4: Anteroposterior (A) and lateral (B) radiographs of a patella fracture involving significant comminution of the inferior pole treated with partial patellectomy. The remaining superior portion is seen on postoperative imaging (C and D).

union. Pin tract infection and/or wire site irritation occurred in 12% but resolved after device removal without further surgical procedures. Articular surface incongruity of 2 mm or greater was seen in 11% of patients, all of whom had radiographic evidence of arthritis at 18 months postoperatively. No secondary surgical procedures were required. The authors concluded that their compression external fixation system was a safe and effective method for treating patella fractures, especially in cases with a poor soft tissue envelope, in salvage situations, and in locations with limited resources.14

Combination Fixation

Some severely comminuted patella fractures may not be amenable to tension band, external fixation, or cerclage wire fixation. To date, little has been published in the literature about the use of combination fixation to include plate and screw osteosynthesis with or without cerclage or tension band fixation. The current authors have found that combining treatment strategies in the setting of significant comminution, especially for high-demand patients in whom partial patellectomy may produce a poor outcome, leads to favorable results (Figure 5).



Figure 5: Anteroposterior (A) and lateral (B) radiographs of a comminuted patella fracture. A medial parapatellar arthrotomy was performed demonstrating multiple fragments (C). The joint surface was reconstructed (D and E) with multiple 2.0-mm screws and a tension band construct. Anteroposterior radiograph (F) showing the final construct.

Total Patellectomy

Total patellectomy is primarily of historical interest and is rarely performed, being reserved for instances of substantial bone loss or as a salvage procedure.^{1,2} Complete patellectomy eliminates the mechanical advantage provided by the patella to the extensor mechanism and results in a 49% reduction in knee extension strength.7,36 A modified technique described by Günal et al40 using a vastus medialis advancement resulted in less pain, less activity limitation, better quadriceps strength, improved cosmesis, and better functional performance than patellectomy alone. However, the authors stated that the patella should be preserved if possible.

OPEN FRACTURES

The patella remains subcutaneous throughout its length, and patella fractures are open 6% to 13% of the time.^{2,41} Open patella fractures tend to result from higher energy mechanisms than closed patella fractures, with motor vehicle accidents causing 94% of open patella fractures and falls causing 62% of closed patella fractures in the same study.⁴¹ Additionally, associated injuries occur in 81% of open fractures compared with 31% of closed patella fractures.41

Treatment of open patella fractures should follow the same principles as treatment of all open fractures: timely, appropriate antibiotics followed by urgent irrigation and





Ε

thorough debridement with definitive fixation and wound closure as soon as possible.2,41

Outcomes following open patella fractures are typically inferior to those following closed patella fractures. However, 65% to 77% good to excellent results have been obtained. Secondary procedures are more common in open patella fractures (up to 65%), and delayed wound coverage/closure is associated with increased risk of deep infection.41-43

COMPLICATIONS

Patient factors have a direct effect on outcomes after surgical treatment of patella fractures; history of a cerebrovascular accident has been found to induce a 6-fold increased risk of infection and a nearly 15-fold increased risk of nonunion.44 Diabetic patients have more than an 8 times increased likelihood of reoperation for all causes.44

Symptomatic hardware, especially in patients treated with a tension band, is common and may occur in up to 60% of patients, often resulting in the need for hardware removal.2,25 Hardware failure occurs in 8% to 22% of patients, most commonly when Kirschner wires are used, and both local and distant hardware migration has been described.45-47 Higher fixation failure rates have been found with increasing patient age and use of Kirschner wires with or without tension band fixation. Increasing duration of follow-up is associated with reoperation and hardware removal, indicating that patellar fixation implants may become more noticeable and symptomatic as time increases after surgery.48

After operative treatment, nonunion and delayed union occurs in 2% to 12.5% of patients and the infection rate ranges from 0 to 5%; both are increased in open fractures. Knee stiffness can best be mitigated by solid fixation and early range of motion. Postoperative radiographic and clinical osteoarthritis are both more common following displaced patella fractures than in the general population and are best minimized by anatomic reduction, solid fixation, and early range of motion.2,21

REHABILITATION

Although numerous clinical protocols have been described. there has been minimal research about the outcomes of specific clinical protocols.^{2,20} Most surgeons recommend gentle early knee range of motion and full weight bearing in a knee brace locked in extension. Flexion is typically allowed to 30° within 2 weeks following surgical fixation with progressive advancement. This may be delayed in cases of extensive comminution or tenuous fixation.2,20

CONCLUSION

Patella fractures represent a broad spectrum of injuries ranging from subtle nondisplaced fractures to open comminuted fractures with significant bone loss. Treatment should be directed to obtaining an anatomic reduction and using a fixation method that maximizes stability while minimizing hardware prominence. Surgeons should select fixation techniques that best address the fracture pattern being treated, as there is little high-quality evidence comparing treatment methods. Despite all of the advances in surgical treatment options, functional impairment, pain, and decreased quadriceps strength and endurance persist to 12 months postoperatively and beyond.10 Knee joint mobilization and range of motion as early as fixation stability permits will help to minimize posttraumatic arthritis and allow optimal postoperative recovery.

REFERENCES

- Boström A. Fracture of the patella: a study of 422 patellar fractures. Acta Orthop Scand Suppl. 1972; 143:1-80.
- Melvin JS, Mehta S. Patellar fractures in adults. J Am Acad Orthop Surg. 2011; 19(4):198-207.
- 3. Kaufer H. Mechanical function of the patella. *J Bone Joint Surg Am.* 1971; 53(8):1551-1560.
- Eckstein F, Müller-Gerbl M, Putz R. Distribution of subchondral bone density and cartilage thickness in the human patella. J Anat. 1992; 180(pt 3):425-433.
- Hung LK, Chan KM, Chow YN, Leung PC. Fractured patella: operative treatment using the tension band principle. *Injury*. 1985; 16(5):343-347.
- Neumann HS, Winckler S, Strobel M. Long-term results of surgical management of patellar fractures [in German]. Unfallchirurg. 1993; 96(6):305-310.
- Carpenter JE, Kasman R, Matthews LS. Fractures of the patella. *Instr Course Lect.* 1994; 43:97-108.
- Böstman O, Kiviluoto O, Nirhamo J. Comminuted displaced fractures of the patella. *Injury*. 1981; 13(3):196-202.
- Tandogan RN, Demirors H, Tuncay CI, Cesur N, Hersekli M. Arthroscopic-assisted percutaneous screw fixation of select patellar fractures. *Arthroscopy*. 2002; 18(2):156-162.
- Lazaro LE, Wellman DS, Sauro G, et al. Outcomes after operative fixation of complete articular patellar fractures: assessment of functional impairment. *J Bone Joint Surg Am.* 2013; 95(14):e96.
- LeBrun CT, Langford JR, Sagi HC. Functional outcomes after operatively treated patella fractures. J Orthop Trauma. 2012; 26(7):422-426.
- Lazaro LE, Wellman DS, Pardee NC, et al. Effect of computerized tomography on classification and treatment plan for patellar fractures. *J Orthop Trauma*. 2013; 27(6):336-344.
- 13. Pritchett JW. Nonoperative treatment of widely displaced

patella fractures. *Am J Knee Surg.* 1997; 10(3):145-148.

- Wardak MI, Siawash AR, Hayda R. Fixation of patella fractures with a minimally invasive tensioned wire method: compressive external fixation. J Trauma Acute Care Surg. 2012; 72(5):1393-1398.
- Lee YM, Chung KJ, Hwang JH, Kim HK, Yoon YH. Results of tension band wiring and additional circumferential wiring in treatment of comminuted patella fracture. *J Korean Fract Soc.* 2014; 27(3):206-212.
- Nummi J. Operative treatment of patellar fractures. Acta Orthop Scand. 1971; 42(5):437-438.
- Sayum Filho J, Lenza M, Teixeira de Carvalho R, Pires OG, Cohen M, Belloti JC. Interventions for treating fractures of the patella in adults. *Cochrane Database Syst Rev.* 2015; 2:CD009651.
- Müller ME, Allgöwer M, Schneider R, Willeneger H. Manual of Internal Fixation: Techniques Recommended by the AO Group. Berlin, Germany: Springer; 1979.
- Gardner MJ, Griffith MH, Lawrence BD, Lorich DG. Complete exposure of the articular surface for fixation of patellar fractures. *J Orthop Trauma*. 2005; 19:118-123.
- 20. Rodrìguez-Merchán E. *Traumatic Injuries of the Knee*. New York, NY: Springer; 2013.
- 21. Berg EE. Open reduction internal fixation of displaced transverse patella fractures with figure-eight wiring through parallel cannulated compression screws. *J Orthop Trauma*. 1997; 11(8):573-576.
- 22. Tian Y, Zhou F, Ji H, Zhang Z, Guo Y. Cannulated screw and cable are superior to modified tension band in the treatment of transverse patella fractures. *Clin Orthop Relat Res.* 2011; 469(12):3429-3435.
- Carpenter JE, Kasman RA, Patel N, Lee ML, Goldstein SA. Biomechanical evaluation of current patella fracture fixation techniques. *J Orthop Trauma*. 1997; 11(5):351-356.
- 24. Gosal HS, Singh P, Field RE. Clinical experience of patellar

fracture fixation using metal wire or non-absorbable polyester: a study of 37 cases. *Injury*. 2001; 32(2):129-135.

- Yang L, Yueping O, Wen Y. Management of displaced comminuted patellar fracture with titanium cable cerclage. *Knee*. 2010; 17(4):283-286.
- Matsuo T, Watari T, Naito K, Mogami A, Kaneko K, Obayashi O. Percutaneous cerclage wiring for the surgical treatment of displaced patella fractures. *Strategies Trauma Limb Reconstr.* 2014; 9(1):19-23.
- 27. Taylor BC, Mehta S, Castaneda J, French BG, Blanchard C. Plating of patella fractures: techniques and outcomes. *J Orthop Trauma*. 2014; 28(9):e231-e235.
- Thelen S, Schneppendahl J, Jopen E, et al. Biomechanical cadaver testing of a fixed-angle plate in comparison to tension wiring and screw fixation in transverse patella fractures. *Injury*. 2012; 43(8):1290-1295.
- Banks KE, Ambrose CG, Wheeless JS, Tissue CM, Sen M. An alternative patellar fracture fixation: a biomechanical study. *J Orthop Trauma*. 2013; 27(6):345-351.
- Wurm S, Augat P, Bühren V. Biomechanical assessment of locked plating for the fixation of patella fractures. *J Orthop Trauma*. 2015. Epub ahead of print.
- 31. Dickens AJ, Salas C, Rise L, et al. Titanium mesh as a lowprofile alternative for tensionband augmentation in patella fracture fixation: a biomechanical study. *Injury*. 2015. Epub ahead of print.
- 32. Wang CX, Tan L, Qi BC, et al. A retrospective comparison of the modified tension band technique and the parallel titanium cannulated lag screw technique in transverse patella fracture. *Chin J Traumatol.* 2014; 17(4):208-213.
- Kastelec M, Veselko M. Inferior patellar pole avulsion fractures: osteosynthesis compared with pole resection. J Bone Joint Surg Am. 2004; 86(4):696-701.
- 34. Yang KH, Byun YS. Separate vertical wiring for the fixation of comminuted fractures of

the inferior pole of the patella. J Bone Joint Surg Br. 2003; 85(8):1155-1160.

- 35. Egol K, Howard D, Monroy A, Crespo A, Tejwani N, Davidovitch R. Patella fracture fixation with suture and wire: you reap what you sew. *Iowa Orthop J*. 2014; 34:63-67.
- Sutton FS Jr, Thompson CH, Lipke J, Kettelkamp DB. The effect of patellectomy on knee function. *J Bone Joint Surg Am.* 1976; 58(4):537-540.
- 37. Bonnaig NS, Casstevens C, Archdeacon MT, et al. Fix it or discard it? A retrospective analysis of functional outcomes after surgically treated patella fractures comparing ORIF with partial patellectomy. J Orthop Trauma. 2015; 29(2):80-84.
- Luna-Pizarro D, Amato D, Arellano F, Hernández A, López-Rojas P. Comparison of a technique using a new percutaneous osteosynthesis device with conventional open surgery for displaced patella fractures in a randomized controlled trial. J Orthop Trauma. 2006; 20(8):529-535.
- 39. Mao N, Ni H, Ding W, et al. Surgical treatment of transverse patella fractures by the cable pin system with a minimally invasive technique. J Trauma Acute Care Surg. 2012; 72(4):1056-1061.
- 40. Günal I, Taymaz A, Köse N, Göktürk E, Seber S. Patellectomy with vastus medialis obliquus advancement for comminuted patellar fractures: a prospective randomised trial.

J Bone Joint Surg Br. 1997; 79(1):13-16.

- Anand S, Hahnel JC, Giannoudis PV. Open patellar fractures: high energy injuries with a poor outcome? *Injury*. 2008; 39(4):480-484.
- Torchia ME, Lewallen DG. Open fractures of the patella. J Orthop Trauma. 1996; 10(6):403-409.
- Catalano JB, Iannacone WM, Marczyk S, et al. Open fractures of the patella: long-term functional outcome. *J Trauma*. 1995; 39(3):439-444.
- 44. Kadar A, Sherman H, Glazer Y, Katz E, Steinberg EL. Predictors for nonunion, reoperation and infection after surgical fixation of patellar fracture. J Orthop Sci. 2015; 20(1):168-173.

- 45. Smith ST, Cramer KE, Karges DE, Watson JT, Moed BR. Early complications in the operative treatment of patella fractures. J Orthop Trauma. 1997; 11(3):183-187.
- 46. Choi HR, Min KD, Choi SW, Lee BI. Migration to the popliteal fossa of broken wires from a fixed patellar fracture. *Knee*. 2008; 15(6):491-493.
- Biddau F, Fioriti M, Benelli G. Migration of a broken cerclage wire from the patella into the heart: a case report. *J Bone Joint Surg Am.* 2006; 88(9):2057-2059.
- Miller MA, Liu W, Zurakowski D, Smith RM, Harris MB, Vrahas MS. Factors predicting failure of patella fixation. J Trauma Acute Care Surg. 2012; 72(4):1051-1055.